

Compost Tea and Its impact On Plant Diseases

by Mario Lanthier

Is compost tea a useful tool to fight plant diseases on organic farms? The answer is "maybe", based on scientific research published recently.

Reports available for many years indicated that foliage sprays of non-aerated compost tea can control rose powdery mildew ¹ and grape powdery mildew ². A thorough review of the topic was published in 2002 by researchers at Oregon State University ³. The survey of 47 studies indicated effective control of mould (*Botrytis cinerea*), tomato late blight (*Phytophthora infestans*), apple scab (*Venturia inaequalis*) and grape powdery mildew (*Uncinula necator*). Persons interested can find a copy of this article on-line.

More recently, aerated compost tea has been promoted as an effective tool to control rose powdery mildew ⁴ as well as grape powdery mildew, leaf anthracnose, peach leaf curl and cherry brown rot ⁵. The claim has been disputed by scientists who failed to control powdery mildew of grapes and apples ⁶.

Compost tea to control powdery mildew

In 2006, the Rodale Institute in Pennsylvania reported on a \$150,000, 2-year study that examined applications of aerated compost tea for control of various leaf diseases in field-grown pumpkins, grapes and potatoes. The recipe for aerated compost tea was reviewed by Soil FoodWeb Inc. in New York ⁷.

Results were erratic for powdery mildew. In pumpkins, there was no impact in 2003 but 80% disease reduction in 2004. In Chardonnay grapes, regular applications of compost tea resulted in 50% decrease in disease incidence on grape clusters in 2003 but had no impact in 2004. The researchers concluded that "compost tea alone does not seem to be a viable tool to suppress powdery mildew on pumpkins" and recommended disease management based on the selection of disease-tolerant grape cultivars.

"The best part of compost tea is the compost" commented Paul Hepperly, the team manager for the Rodale Institute project. "In potatoes, we found advantages to compost tea only when soil did not receive compost." ⁸



Shown is Elise L. Brun in the orchard of a cooperating grower near Cawston, B.C. Aerated compost tea was sprayed on the foliage at weekly intervals from April to July in an attempt to prevent powdery mildew. Results indicated a significant impact from the commercial fungicide Kumulus, but inconsistent control with compost tea.

In 2005, the Organic Food and Farming Education and Research program of Ohio State University reported on a 1-year study that examined various products in organic squash production. Aerated compost tea was prepared with the commercial brewer "SoilSoup". Powdery mildew was significantly lower following sulphur treatments, but there was no reduction from compost tea applications ⁹.

The claim that aerated compost tea can prevent powdery mildew remains anecdotal. For powdery mildew, this author agrees with a comment posted recently on a web discussion group: "As hard as I search, I still have not found an article on disease suppression supporting aerated compost tea." ¹⁰

Compost tea to control other diseases

In 2004, researchers from the U.S. Department of Agriculture in Oregon reported that a drench application of compost tea was effective to suppress damping-off of cucumber (caused by *Pythium ultimum*) grown in soiless greenhouse media. However, the results were inconsistent when aerated tea was prepared without additives, and results were also inconsistent when non-aerated tea was prepared with or without additives ¹¹.

...continued on page 8

...continued from page 7



Botrytis mold growing on a plant part
This disease is common in many agriculture crops such as greenhouses, vegetables, tree fruits and berries. It typically infects weak or dying plant parts and later spreads to healthy tissue. Scientific research indicates that compost tea is effective in prevention of Botrytis mold.

The most consistent disease suppression was obtained from aerated compost tea fermented with kelp and humic acids.

Kelp and humic acids alone did not suppress damping-off, but triggered disease suppression when added to any of three different types of compost (yard trimmings, vermicompost, or a proprietary blend of tea compost). Diluting the finished tea with water, or imposing heat treatment (to kill live micro-organisms) significantly reduced suppression, indicating the impact was related to microbes but not nutrients.

The authors suggest an *aerated* compost tea becomes "disease suppressive" when made without molasses and containing a bacterial population of 7 log₁₀ Colony Forming Units (CFU) / ml of solution.

In 2006, the same researchers reported on the impact of compost tea recipes against grey mould (caused by *Botrytis cinerea*) of geranium. For *non-aerated* compost tea, only 31% of teas tested suppressed grey mould. The most consistent and significant disease suppression came from teas made of composted chicken manure or composted yard waste. Increasing fermentation time from 7 to 14 days significantly reduced disease. Adding nutrients or stirring during fermentation did not help with disease suppression ¹².

In this project, the *aerated* compost tea was prepared with commercial brewers from "Growing Solutions, Inc." or "Soil Soup, Inc.". Only 17% of teas tested suppressed grey mould, but 67% of tea batches significantly reduced disease when

made in the presence of kelp and humic acid. Addition of an adjuvant (such as spreader or sticker) prior to application significantly reduced disease levels compared to aerated tea or adjuvant alone, possibly from increased attachment to the leaf surface.

In 2003, scientists at Ohio State University prepared *non-aerated* compost tea from either composted cow manure, composted pine bark or organic farm compost. The water extract was sprayed on foliage of tomato plants and tested against bacterial spot (caused by *Xanthomonas vesicatoria*) ¹³.

In laboratory assays with transplants, a foliage spray 24 hours before disease infection resulted in a significant reduction in bacterial spot severity. There was no difference in efficacy between teas prepared from "younger" or "older" compost (curing of 3 to 16 months) or compost to water ratios of 1:1, 1:3 and 1:5. In field production, there was no difference between weekly and biweekly sprays of compost tea. Efficacy was marginal and considerably less than standard copper hydroxide sprays at weekly intervals.

Finally, in a recent study at Penn State University for management of apple scab (caused by *Venturia inaequalis*), a post-harvest foliar application of *aerated* compost tea significantly reduced spring ascospore production compared to the water check ¹⁴.

Home-size compost tea recipe

From W.F. Brinton at Woods End Laboratories Inc., Maine. See the website <http://www.woods-end.org/compostteas.html>.

- 1) Use well-aged compost, at least 4 months old.
- 2) Put in a large pail or barrel outdoors between 15 and 20°C.
- 2) Add water, 1 part compost to 5 to 8 parts water (i.e. 1 cup of compost to 5 cups of water).
- 3) Stir daily for five days. The strong smell should slowly dissipate.
- 4) On the fifth day, pour through a sieve or a cheese cloth.
- 5) Spray on plants in periods of disease outbreak, or drench the soil at the base of the plant.
- 6) Do not spray edible plant parts to be harvested in the following 2 to 3 weeks.

The bottom line

There is strong scientific evidence that aerated compost tea can prevent a number of plant diseases such as damping off and Botrytis mold, but it is not effective against powdery mildew.

For best results, start the tea with high quality compost, for example a mixture of farm compost and vermicompost. Add kelp and humic acid during brewing. The procedure aims to extract and grow beneficial live microorganisms found in the compost.

Results are not as good with non-aerated compost tea. Again, using high quality compost is critical to obtain success. The procedure aims to extract antibiotic compounds found in the compost.

Until further research anchors the rates and application techniques, growers should use compost tea as a supplement to sound cultural practices and organic-approved spray applications.

Understanding modes of action

Different types of compost tea behave in different ways when applied to plants. A solid understanding of these modes of action is crucial to make successful use of compost tea against plant diseases.

Preparation of non-aerated compost tea favours the extraction of antibiotic compounds that play an important role in suppression of plant pathogens.

One study examined compost tea that inhibited apple scab. When the compost was sterilized before fermentation, the tea was not disease suppressive. However, when the compost was not sterilized but the finished tea was autoclaved, the tea retained its disease suppression properties, indicating the disease suppression properties did *not* come from live microorganisms. The inhibitory agent was determined to be a low molecular weight, heat stable, non-protein metabolite produced by microorganisms during fermentation ¹⁵. A similar conclusion was reached by another group who compared efficacy of finished tea against autoclaved finished tea

for space and nutrients. For other pathogens, such as powdery mildew and rust, spore germination and plant penetration can occur in the absence of exogenous nutrients, and microbial competition would *not* prevent pathogen growth ¹⁸.

Soil management may offer a long-term prospect for suppression of leaf diseases such as powdery mildew. Specific soil microbes colonizing plant roots can induce plant production of defence-related proteins, resulting in increased resistance to foliage diseases ¹⁹.



Compost tea for organic farming

Shown is Shepard Smith, then an organic vegetable grower in Oregon. He is seen inspecting a commercial machine for aerated compost tea. The white container is a reservoir for water and compost and is equipped with a pump to deliver constant air during the brewing process.

...continued on page 10

Types of compost tea

Non-aerated compost tea describes procedures where compost is mixed with water and left to stand for many days with minimal disturbance. It is also called "compost extract", "compost slurry" or "steepage".

Different methods were recently reviewed²⁰. In the 1980s, a team of German researchers made "watery fermented compost extract" from compost mixed with water in a ratio of 1:5 to 1:8 in an open container, stirred once then allowed to sit for 5 to 8 days. The end product was filtered to remove large debris and the liquid extract used immediately.

The recipe was slightly modified by other researchers, including Elad and Shtienberg in the 1990s (compost:water ratio of 1:5 with a 10-day extraction period²¹) and Al-Dahmani *et al* in 2003 (compost:water ratio of 1:5 placed in a plastic container and stirred twice during a 7-day incubation at 20 to 22°C¹²). Woods End Research Laboratory in Maine recommends fermentation in wooden barrels at 15 to 25°C for 3 to 8 days with 2 or 3 stirs using a rod or rotating bar. The end product is not stirred for 8 hours before filtration (to avoid clogging spray equipment), decanted through a 200-mesh sieve (75-micron) and mixed with a proper wetter / sticker agent to ensure full plant coverage²².

Actively aerated compost tea is a more recent technology where the mixture of compost and water is supplied with active aeration, for example with an aquarium pump. The first "compost tea brewer" was conceived in 1993 and today different models are available for purchase²³.

The high oxygen concentration stimulates population growth of aerobic microbes, which helps disease prevention, nutrient cycling, retention of micro-nutrients, soil structure, and decomposition of plant-toxic materials. By contrast, these beneficial microbes may not survive in *non-aerated* compost tea because of anaerobic conditions⁴.

Basic procedures for *aerated* compost tea are readily available²⁴. Instructions for the construction of a home-made unit are posted on the website of the Pennsylvania Department of Environmental Protection at <http://www.dep.state.pa.us/dep/deputate/airwaste/wm/recycle/Tea/tea1.htm>²⁵.

B.C. research on powdery mildew

Weekly application of *aerated* compost tea is not effective to control powdery mildew in apple trees. That was the conclusion of a research project conducted in 2004 in the Similkameen Valley²⁶.

The trials were placed within newly-planted apple trees near Cawston, B.C. Five treatments replicated four times were distributed in a randomized, replicated block design. Plot size was 10 consecutive trees. Treatments were applied with a hand-held back-pack to the tree foliage during the morning, at weekly intervals between late April and early July (ten applications). Actively aerated compost tea was prepared with a commercial brewer using local well water and a package of compost plus additives supplied by the brewer manufacturer. Tea was applied shortly after preparation in a spray solution of 10% dilution. Disease incidence was left to occur from surrounding areas. Rating was done on 800 leaves per treatment²⁷ and results analysed for treatment significance.

Disease incidence was moderate at the first orchard and low at the second orchard. The commercial fungicide Kumulus resulted in a significant reduction in powdery mildew at both sites. Compost tea alone, or spreader alone, resulted in a significant reduction in powdery mildew at the first site, but had no measurable impact at the second site.

Preparation of aerated compost tea

Shown is Sonja Peters with a 5-gallons commercial brewer.

In non-aerated compost tea, compost is added to water in a large container and let to ferment for 7 to 10 days. In aerated compost tea, air is forced into the mixture to stimulate growth of beneficial microorganisms.



Table 1. Mean % leaf surface occupied by powdery mildew on young apple trees following 10 weekly applications from late April to early July 2004 (each result is the mean of 800 ratings) ¹

Treatment	Rate	Orchard # 1 Rating on 30 June	Orchard # 2 Rating on 8 July
Untreated control		7.5	0.20
Grower standard ²	20 gr / 10 L water	5.6 **	0.06 **
Compost tea ³	1 L / 10 L water	6.8 *	0.20
Spreader only ⁴	10 ml / 10 L water	6.5 *	0.19
Compost tea + Spreader	As above	7.3	0.16
Standard error		0.22	0.013
Significance		* signifies $p < 0.05$	** signifies $p < 0.001$
		** signifies $p < 0.001$	

1: Data analysed by Dr. Ben Coleman, Okanagan College, Kalamalka Campus. Variability between group means was examined with F-test ($p < 0.05$). Significance between treatments was determined with pairwise comparison (Tukey's HSD).

2: Commercial fungicide Kumulus containing 90% sulphur

3: Commercial 5-US capacity brewer and ingredients from K.I.S. (Keep It Simple Inc., www.simpli-tea.com)

4: Commercial product "Superflow" from BiozAgri Products, Oliver B.C. (www.raingrow.com)

5: Stars indicate a significant condition between the treatment and untreated, Tukey's HSD

Funding for this project came from the Certified Organic Associations of B.C. (the Organic Sector Development Program) and cooperating organic farmers (J. and G. Dhaliwal, L. and G. Sellmer, J. and R. Mennell, S. and W. Mennell, L. Edwards and B. Mennell).

FOOTNOTES

- 1- Scheuerell S.J. and W.F. Mahaffee. 2000. "Assessing aerated and nonaerated watery fermented compost and *Trichoderma harzianum* T-22 for control of powdery mildew (*Spraecotheca pannosa* var. *rosae*) of rose in the Willamette Valley, Oregon". *Phytopathology*. 90s: 69.
- 2- Trankner A. 1992. "Use of agricultural and municipal organic wastes to develop suppressiveness to plant pathogens". In: *Biological control of plant diseases*. Tjamos E.C., G.C. Papavizas and R.J. Cook editors. Plenum Press (New York NY). Pages 35-42.
- 3- Scheuerell S. and W. Mahaffee. 2002. "Compost tea: Principles and prospects for plant disease control". *Compost Science & Utilization*. 10(4): 313-338.
- 4- Ingham E.R. 2005. "The Compost Tea Brewing Manual". Soil Foodweb Incorporated (Corvallis, OR). Fifth Edition. Page 51.
- 5- Rollins C.A. editor. 2004. "The Field Guide to Actively Aerated Compost Tea (AACT)". Soil Foodweb, Inc. (Corvallis, OR) and Nature Technologies, LLC (Sonoma CA). Page 50
- 6- Pscheidt J. and H. Wittig. 1996. "Fruit and Ornamental Disease Management Testing Program". Plant Pathology Extension. Oregon State University (Corvallis, OR).
- 7- Hepperly P. 2006. "Compost tea for disease management in horticultural crops". Sustainable Agriculture Research and Education (SARE) Project Number LNE03-181. Available at http://www.sare.org/reporting/report_viewer.asp.
- 8- "A cuppa compost". *Organic Gardening*. April/May 2006 issue. Page 14.
- 9- Miller S. *et al.* 2005. "Disease management for organic vine crops". In: *Progress Report of the Organic Food and Farming Education and Research Program*. Ohio Agricultural Research and Development Center. Pages 46-47.
- 10- Posting by Matthew Ryan, 25 October 2004. Web discussion group http://groups.yahoo.com/group/compost_tea
- 11- Scheuerell S.J. and W.F. Mahaffee. 2004. "Compost tea as a container medium drench for suppressing seedling damping-off caused by *Pythium ultimum*". *Phytopathology*. 94: 1156-1163.
- 12- Scheuerell S.J. and W.F. Mahaffee. 2006. "Variability associated with suppression of gray mold (*Botrytis cinerea*) on geranium by foliar applications of nonaerated and aerated compost tea". *Plant Disease*. 90: 1201-1208.
- 13- Al-Dahmani J.H. *et al.* 2003. "Suppression of bacterial spot of tomato with foliar sprays of compost extracts under greenhouse and field conditions". *Plant Disease*. 87: 913-919.
- 14- Travis J.W. *et al.* 2006. "Abstract book". Page 115e. American Phytopathological Society annual meeting.
- 15- Cronin *et al.*, 1996. In: Scheuerell S. and W. Mahaffee. 2002. "Compost tea: principles and prospects for plant disease control". *Compost Science & Utilization*. 10(4): 313-338.
- 16- Zhang W. *et al.* 1998. "Compost and compost water extract-induced systemic acquired resistance in cucumber and Arabidopsis". *Phytopathology*. 88: 450-455.
- 17- Hoitink H.A., A.G. Stone, D.Y. Han. 1997. "Suppression of plant diseases by composts". *HortScience*, 32(2): 184-187.
- 18- Elad Y., N.E. Malathrakis and A.J. Dik. 1996. "Biological control of *Botrytis*-incited diseases and powdery mildews in greenhouse crops". *Crop Protection*. 15(3): 229-240.
- 19- Hoitink H.A.J. *et al.* 2006. "Systemic resistance induced by *Trichoderma* spp.: Interactions between the host, the pathogen, the biocontrol agent, and soil organic matter quality". *Phytopathology*. 96:186-189.
- 20- Quarles W. 2001. "Compost tea for organic farming and gardening". *The IPM Practitioner*. XXIII(9): 1-8.
- 21- Elad Y. and D. Shtienberg. 1994. "Effect of compost water extracts on grey mould (*Botrytis cinerea*)". *Crop Protection*. 13(2): 109-114.
- 22- Brinton W.F., A. Trankner and M. Droffner. 1996. "Investigations into liquid compost extracts". *BioCycle*. 37(11): 68-70.
- 23- Information on commercial brewers is various websites, including http://www.soilfoodweb.com/03_about_us/approach_pg/c_03a_aerated_tea.html and <http://www.jollyfarmer.com/compost%20tea%20brewers.htm>.
- 24- Ingham E. 2002. "A rated aerated compost tea". *EcoFarm & Garden*. Summer issue., 18-20.
- 25- Pennsylvania Department of Environmental Protection. "Compost tea as easy as 1 2 3". Retrieved January 2007 at <http://www.dep.state.pa.us/dep/deputate/airwaste/wm/recycle/Tea/tea1.htm>.
- 26- Lanthier M. and S. Peters. 2006. "Compost tea: Testing the field application of actively aerated compost tea for prevention of powdery mildew and improvement of soil microbial community in fruit tree production in British Columbia". Organic Sector Development Program Project #I0933. Certified Organic Associations of British Columbia. Vernon BC. Summary report on objective 2, Appendix 1, Appendix 2.
- 27- Belanger R.R. *et al.* 2002. "The powdery mildews: A comprehensive treatise". American Phytopathological Society Press (St. Paul, MI). Pages 94-95.